

# Targeted Copy Number Variations Profiling of Non Muscle Invasive Bladder Urothelial Carcinoma Using BCA-1 Test on Urines Predicts High Grade Tumors

Lunelli L<sup>1,4</sup>, Léon P<sup>1,2</sup>, Cancel Tassin G<sup>1,2,3</sup>, Sighar K<sup>6</sup>, Compérat E<sup>1</sup>, Hugonin S<sup>5</sup>, Audouin M<sup>1,4</sup>, Lacave R<sup>5</sup> and Cussenot O<sup>1,3,4\*</sup>

<sup>1</sup>Department of Urology, Sorbonne University - UPMC, France

<sup>2</sup>Academic Department of Urology, CHU Reims, France

<sup>3</sup>CeRePP - Pathologies Prostatiques - Centre de Recherche, France

<sup>4</sup>Academic Department of Urology, UPMC Sorbonne University, France

<sup>5</sup>Department of Tumor Biology, Hospital Tenon, France

<sup>6</sup>Arraygenomics, Voisins le Bretonneux, France

## ARTICLE INFO

### Article history:

Received: 08 August 2017

Accepted: 02 October 2017

Published: 19 October 2017

### Keywords:

NMI- TCC:

CNV:

UUT-TCC:

BCA1:

Cystoscopy

**Copyright:** © 2017 Cussenot O

Urol Res Ther J

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Citation this article:** Cussenot O, Lunelli L, Léon P, Cancel Tassin G, Sighar K et al. Targeted Copy Number Variations Profiling of Non Muscle Invasive Bladder Urothelial Carcinoma Using BCA-1 Test on Urines Predicts High Grade Tumors. Urol Res Ther J. 2017; 1(1):113.

### Correspondence:

Dr. Cussenot O,  
Department of Urology, Hospital  
Tenon, AP-HP, 4 rue de la Chine,  
Paris, F-75020, France,  
Email: olivier.cussenot@aphp.fr

## ABSTRACT

**Introduction:** Urothelial carcinomas are the fourth leading cause of cancer in humans. Their incidence has increased by more than 50% in 25 years. Non Muscle Invasive Bladder Urothelial Carcinoma (NMI-TCC) represent 70% of newly diagnosed cases. Bladder NMI-TCC with high grades or pT1 stages require a close active surveillance regarding their significant risk of recurrence and progression to an invasive stage. From results obtained using DNA markers, it has been recognised that FGFR3 mutations were most associated to low grade and pTa NMI-TCC, while numerous CNV were associated to genetically instable tumors with high grade or pT1/pTis stage. The BCA-1 test provides a targeted copy number variations (CNV) profiling. So, we evaluated the predictive value of the FGFR3 mutations and BCA-1 test on urines in order to assess the tumoral grade and the stage of NMI-TCC.

**Method:** Urinary DNA was extracted from 50 patients with bladder NMI-TCC, with negative or non-informative urine cytology. The BCA1 test and identification of FGFR3 mutations (S249C; Y375C; G372C) was performed on urine samples. The molecular results were correlated with the tumoral pathological grade and stage.

**Results:** Each CNV locus was correlated to grade and stage. Gains at 8q, 11q and deletions at 9p 9q loci were the most informative CNV for high grade and pT1 prediction. A Bayesian model from molecular markers using urines CNV profiling predicts high grade and pT1 stage, with AUC respectively at 92% and 93%.

**Conclusion:** Our results showed that targeted CNV profiling using BCA-1 test on urines could be used as biomarkers during clinical management of bladder NMI-TCC, as a reliable predictor of tumors with high risk of recurrence and progression.

### Introduction

Nearly 180,000 new cases of bladder cancer are diagnosed each year in the European Union, with a ratio male/female of 2/3. The main risk factor is due to cigarettes smoking habits. At the time of diagnosis, around 70% of bladder tumors are Non-Muscle Invasive Transitional Cell Carcinoma (NMI-TCC) [1].

The stages and grades distribution of the bladder NMI-TCC newly diagnosed are respectively about 60% as pTa, 40% as pT1 and <5% pTis. Half of the distribution of bladder NMI-TCC is tumor with high grade. Urinary cytology usually performed in clinical practice has a low sensitivity for low- grade tumors and require experimented pathologist to perform this examination [2]. It is currently recognized than cytology has a sensitivity of 69–92% for high-grade tumors but this does not make it a good monitoring test [2]. One common feature to many tumors is the presence of chromosomal anomalies which can be detected by studying DNA markers [3]. This particular characteristic of the tumoral genome was first used in bladder cancers to detect tumoral cells by studying DNA extracted from patient urine with the help of microsatellite markers [4,5]. Several teams, including us, have since reported results where this technique has been used to diagnose or monitor such tumors [6,7]. We and other have, also, previously shown than molecular urine test using a panel of genomic markers identified with a high sensitivity and specificity (over 85%), recurrence of bladder NMI-TCC [4,8]. Since then, BCA1-test based on a targeted CNV panel (60,000 probes with 50-60 base pairs) performed on urine shown no false positives in a population of patients with normal cystoscopy, normal cytology and no history of bladder cancer, and showed a sensitivity of 95% for detecting positive cystoscopies in patients monitored for bladder cancer [9].

We have now extended our study to the prospective analysis of patients with superficial bladder cancer in order to identify the best CNV markers able to predict tumors grade and/or stage.

**Methods**

A nested cohort of 50 patients with a bladder tumor was included in this study. The criteria for inclusion were firstly an indication of endoscopic bladder resection in the context of bladder NMI- TCC (<3cm) monitoring or recurrence; secondly, a negative or non-informative cytology; thirdly, no sign of **Upper Urinary Tract Tumors** (UUT-TCC) on CT-scan. All the patients gave their informed consent. The clinical data collected included

age, sex, age on diagnosis, upper urinary tract tumor history, smoker status (active, former or non-smoker). The following anatomo-pathological data about the bladder tumor were also collected: the stage and grade, and the mutational status of the FGFR3 gene for S249C; Y375C; G372C mutations [10].

**Table 1:** Characteristics of the 50 patients included.

	Number (%)
Men	44 (88%)
Women	6 (12%)
Median age on diagnosis (range)	67 (35 – 90)
Non-smoker	12 (24%)
Former-smoker	29 (58%)
Active Smoker	9 (18%)
UUT-TCC history	5 (10%)
pTa	36 (72%)
pT1	14 (28%)
High-grade	22 (44%)
FGFR3 Mutation (S249C, Y375C or G372C)	11 (22%)

Bladder NMI-TCC (<3cm), negative or non informative cytology, no sign UUT-TCC on CT- scan


**Collection of Urine Samples**

Patients include in the cohort, have sign a consent according to their participation to the protocols Protocole 2011/24 NICB : (IRB 00003835 Institutional Review Board) Comité de Protection des Personnes Ile de France IV - Saint Louis, 1, Avenue Claude Vellefaux 75010 PARIS). Urine samples (50 ml) were collected before each intervention. After centrifuging the urine, the QIAmp DNA Blood Mini kit (Qiagen) was used to extract the urinary DNA from the pellet obtained, complying with the supplier’s recommendations. After validation of the quantity and quality of the extracted DNA, it was marked then hybridized on the BCA-Oligo array following the protocol described in Larré et al. [8].

Briefly, the BioPrime kit (Life Technologies Invitrogen) was used in accordance with the supplier’s recommendations to mark the tested DNA (with Cyanine 5) and the reference DNA (with Cyanine 3) [11,12]. After double purification, the first with the Nucleospin Extract II kit (Macherey-Nagel) and the second with the Purelink® kit (Life Technologies), the two DNAs were mixed in equal quantity. The mixture was then incubated for 3 minutes at



After washing with the Wash Buffer washing solution (Agilent Technologies), the array was scanned using the Agilent High Resolution Scanner (Agilent Technologies). WorkBench software (Agilent Technologies) was used to analyze the results.




To interpret the outcome of the BCA-Oligo test, the median Log2 normalized fluorescence value was calculated for each oligonucleotide. An oligonucleotide was considered deleted or amplified if its normalized average Log2 ratio was lower or higher than -0.2 or 0.2 respectively. A chromosomal region was considered amplified or deleted if at least 6 successive oligonucleotides of the region were interpreted as gained or lost. If at least one chromosomal region was amplified or deleted, the test was considered positive for the study.

**Statistics**

Descriptive statistics were performed using (XLS-STAT® 2017.4). Correlations, relative importance and predictive models were performed using naïve and supervised Baeyesian networks analysis (BayesiaLab BayesiaLab® 6.0.8, Laval, France). Values of  $p < 0.05$  were considered statistically significant. [13]

**Discussion**

The management of bladder with a high grade or pT1/pTis stages require specific follow-up regarding the high risk (>50%) of recurrence or disease progression to muscle invasive stages [12,14]. Our study, using BCA-1 test on urine points out chromosomal locus, such as 8q or 11q or 9q with specific CNV related to high grade or pT1 bladder NMI-TCC. So, Zaharieva et al. [15] previously showed that the frequency of gains of the cMYC gene situated in 8q24 increases significantly with the grade and stage (frequency of 6% for grade G1, 12% for G2 and 20% for G3; 11% for pTa to 16% for pT1). In a study of 28 pTa and 28 pT1 tumors, Richter et al. [16] showed that the frequency of losses in 8p and gains in 3p, 8q and 10p was significantly higher in pT1 tumors. In another study of 64 Chinese patients with bladder tumors, the gains in 8q and 11q and the losses in 8p were more frequent in pT1 tumors than in pTa tumors [4].



A study of 46 superficial tumors showed that the losses in 8p and amplifications in 8q, 5q and 10p were more frequent at the pT1 stage [17]. FGFR3 mutation were frequently (40%) associated to low grade disease, but not exclusively. The fact that FGFR3 mutation are inconstant in low grade, enhance the fact negative FGFR3 mutation is not a reliable markers if used alone [10]. These results suggest that the use of CNV to predict the risk of aggressiveness or evolution of bladder NMI-TCC from DNA extracted from urine samples could be easily done. However, our predictive model needs to be confirmed on an independent larger cohort of patients.

### Conclusion

As a complement to cytology, the identification of markers of aggressiveness in a patient can help the clinician to more easily take a decision on adjuvant therapy after resection of the tumor or to devise a personalized therapeutic strategy. Establishing distinct genetic profiles of patients with bladder NMI-TCC could help to improve care in terms of both diagnosis and prognosis assessment.

### Personal interests

Karim Sighar is an employee of Array Genomics, which helped to develop and market the BCA-oligo test (BCA-1). The other authors state that they have no personal interests.

### Results

The characteristics of the population studied were presented in the Table 1. Individual results were presented in (Figure 1). The distribution and the typology (Gains, Deletions, Others) of each targeted locus for CNV profile were shown on (Figure 2). The cartography of the relationship between clinical, pathological and molecular factors is represented on (Figure 3). Significant correlations are link between age and smoking habits (Pearson R: -0.7; p value<0.001), high grade and stage pT1 (Pearson R: 0.7 ; p value<0.001) ; stage pT1 and CNV (Pearson R: 0.5 ; p value: 0.002). The informativeness each target locus for high grade was shown on Figure 4. The more informative (relative importance >0.3) markers were gains at 8q ,11q and deletions at 9q. From this cartography the predictive

value of the Bayesian model give an AUC respectively at 92% and 93% for high grade and pT1.

### References

1. Burger M, Catto JW, Dalbagni G, Grossman HB, Herr H, et al. (2013). Epidemiology and risk factors of urothelial bladder cancer. *Eur Urol.* 63: 234-241.
2. Babjuk M, Böhle A, Burger M, Capoun O, Cohen D, et al. (2017). EAU Guidelines on Non-Muscle-invasive Urothelial Carcinoma of the Bladder: Update 2016. *Eur Urol.* 71: 447-461.
3. Mao L, Lee DJ, Tockman MS, Askin F, Sidransky D, et al. (1994). Microsatellite alterations as clonal markers for the detection of human cancer. *Proc Natl Acad Sci USA.* 91: 9871-9875.
4. Qin SL, Chen XJ, Xu X, Shou JZ, Bi XG, et al. (2006). Detection of chromosomal alterations in bladder transitional cell carcinomas from Northern China by comparative genomic hybridization. *Cancer Lett.* 238: 230-239.
5. Steiner G, Schoenberg MP, Linn JF, Mao L, Sidransky D, et al. (1997). Detection of bladder cancer recurrence by microsatellite analysis of urine. *Nat Med.* 3: 621-624.
6. Mourah S, Cussenot O, Vimont V, Desgrandchamps F, Soliman H, et al. (1998). Assessment of microsatellite instability in urine in the detection of transitional-cell carcinoma of the bladder. *Int J Cancer.* 79: 629-633
7. Van Rhijn BW, Lurkin I, Kirkels WJ, van der Kwast TH, Zwarthoff EC, et al. (2001). Microsatellite analysis -DNA test in urine competes with cystoscopy in follow-up of superficial bladder carcinoma: a phase II trial. *Cancer.* 92: 768-775.
8. Larré S, Camparo P, Comperat E, Traxer O, Roupret M, et al. (2011). Diagnostic, staging and grading of urothelial carcinomas from urine: performance of BCA-1, a mini-array comparative genomic hybridization based test. *Eur Urol.* 59: 250-257.
9. Cussenot O, Sighar K, Mohammed M, Hugonin S, Ondet V, et al. (2014). Detection of specific chromosomal aberrations in urine using BCA-1 (oligo-

CGH-array) enhances diagnostic sensitivity and predicts the aggressiveness of non-muscle-invasive bladder transitional cell carcinoma. *World J Urol.* 32: 551-557.

10. Couffignal C, Desgrandchamps F, Mongiat-Artus P, Ravery V, Ouzaid I, et al. (2015). The Diagnostic and Prognostic Performance of Urinary FGFR3 Mutation Analysis in Bladder Cancer Surveillance: A Prospective Multicenter Study. *Urology.* 86: 1185-1190.

11. Hoglund M, Sall T, Heim S, Heim S, Mitelman F, et al. (2001). Identification of cytogenetic subgroups and karyotypic pathways in transitional cell carcinoma. *Cancer Res.* 61: 8241-8246.

12. Hoque MO, Lee CC, Cairns P, Schoenberg M, Sidransky D, et al. (2003). Genome-wide genetic characterization of bladder cancer: a comparison of high-density single-nucleotide polymorphism arrays and PCR-based microsatellite analysis. *Cancer Res.* 63: 2216-2222.

13. Roupret M, Hupertan V, Yates DR, Comperat E, Catto JW, et al. (2008). A comparison of the performance of microsatellite and methylation urine analysis for predicting the recurrence of urothelial cell carcinoma, and definition of a set of markers by Bayesian network analysis. *BJU Int.* 101: 1448-1453.

14. Comperat E, Varinot J. (2016). Immunochemical and molecular assessment of urothelial neoplasms and aspects of the 2016 World Health Organization classification. *Histopathology.* 717-726.

15. Zaharieva B, Simon R, Ruiz C, Oeggerli M, Mihatsch MJ, et al. (2005). High-throughput tissue microarray analysis of CMYC amplification in urinary bladder cancer. *Int J Cancer.* 117: 952- 956.

16. Richter J, Jiang F, Görög JP, Sartorius G, Egenter C, et al. (1997). Marked genetic differences between stage pTa and stage pT1 papillary bladder cancer detected by comparative genomic hybridization. *Cancer Res.* 57: 2860-2864.

17. Simon R, Burger H, Brinkschmidt C, Bocker W, Hertle L, et al. (1998). Chromosomal aberrations associated with invasion in papillary superficial bladder cancer. 185: 345-351.